

# A Landesman-Lazer type condition for asymptotically linear second order equations with a singularity

Alessandro Fonda and Maurizio Garrione\*

## Abstract

We consider the  $T$ -periodic problem

$$\begin{cases} x'' + g(t, x) = 0 \\ x(0) = x(T), x'(0) = x'(T), \end{cases}$$

where  $g : [0, T] \times ]0, +\infty[ \rightarrow \mathbb{R}$  exhibits a singularity of a repulsive type at the origin, and an asymptotically linear behavior at infinity. In particular, for  $x$  large,  $g(t, x)$  is controlled from both sides by two consecutive asymptotes of the  $T$ -periodic Dancer-Fučik spectrum, with possible equality on one side. By means of a suitable Landesman-Lazer type condition, we prove the existence of a solution.

## 1 Introduction

In [6], del Pino, Manásevich and Montero considered the problem of finding  $T$ -periodic positive solutions of the equation

$$x'' + g(t, x) = 0, \quad (1)$$

where  $g : \mathbb{R} \times ]0, +\infty[ \rightarrow \mathbb{R}$  is continuous,  $T$ -periodic in its first variable and has a singularity of a repulsive type at the origin. They proved the following result.

**Theorem 1.1.** (*[6, Theorem 1]*) *Assume that there are an integer  $N$  and two constants  $\hat{\mu}, \tilde{\mu}$  such that*

$$\left(\frac{N\pi}{T}\right)^2 < \hat{\mu} \leq \liminf_{x \rightarrow +\infty} \frac{g(t, x)}{x} \leq \limsup_{x \rightarrow +\infty} \frac{g(t, x)}{x} \leq \tilde{\mu} < \left(\frac{(N+1)\pi}{T}\right)^2, \quad (2)$$

*uniformly for every  $t \in [0, T]$ . Moreover, suppose that there are positive constants  $c', c'', \delta$  and  $\nu \geq 1$ , such that*

$$\frac{c'}{x^\nu} \leq -g(t, x) \leq \frac{c''}{x}, \quad \text{for every } t \in [0, T] \text{ and every } x \in ]0, \delta]. \quad (3)$$

*Then, equation (1) has a  $T$ -periodic solution.*

---

\*The authors wish to acknowledge the support of G.N.A.M.P.A., in the frame of the project “Soluzioni periodiche di alcune classi di equazioni differenziali ordinarie”.

Assumption (2) seems to be a kind of nonresonance condition with respect to the set

$$\Sigma = \left\{ \left( \frac{k\pi}{T} \right)^2, k \in \mathbb{N} \right\},$$

whose elements correspond to the heights of the asymptotes of the curves in the Dancer-Fučik spectrum (see, e.g., [10]). They also coincide with the eigenvalues of the associated Dirichlet problem on the interval  $[0, T]$ . As a direct consequence of Theorem 1.1, the equation

$$x'' - \frac{1}{x^\nu} + \beta x = e(t), \quad (4)$$

where  $\nu \geq 1$  and  $e(t)$  is continuous and  $T$ -periodic, has a  $T$ -periodic solution, provided that  $\beta > 0$  satisfies  $\beta \notin \Sigma$ .

Let us mention that the study of the existence of periodic solutions for equations with a singularity, like (1), was first considered by Lazer and Solimini in [14], and later investigated by many authors, see e.g., [2, 4, 5, 7, 8, 9, 11, 17, 19, 20, 21, 22, 23].

A few lines below the statement of their main theorem, del Pino, Manásevich and Montero raised the problem of finding sufficient conditions on  $e(t)$  in order to ensure the existence of a  $T$ -periodic solution for equation (4) in the case when  $\beta$  belongs to the set  $\Sigma$ . As a matter of fact, this seems to be a delicate problem for the following reason. As already remarked in [6] (see also [1, 16]), if  $\nu = 3$ , in the phase plane there is an isochronous center for the Steen's equation (see [18])

$$x'' - \frac{1}{x^3} + \beta x = 0, \quad (5)$$

for any choice of  $\beta > 0$ , since all the positive solutions have minimal period equal to  $\pi/\sqrt{\beta}$ . This fact is crucial when we deal with the  $T$ -periodic problem associated with (5), in the case when  $\beta \in \Sigma$ . Indeed, by a phase plane analysis it was shown in [16] that, due to isochronicity, if one considers a forcing made by a  $T$ -periodic chain of Dirac deltas, then all the solutions of the forced equation are unbounded, both in the past and in the future. The same type of situation arises for the equation

$$x'' - \frac{1}{x^3} + \beta x = \varepsilon \sin(\sqrt{\beta}t),$$

for  $|\varepsilon|$  sufficiently small, as proved in [4, Theorem 3].

However, in the case  $\nu = 3$ , defining the  $(\pi/\sqrt{\beta})$ -periodic function

$$\Phi(\theta) = \int_0^T e(t) |\sin(\sqrt{\beta}(t + \theta))| dt,$$

it was shown in [4] that if  $\Phi$  has only simple zeros, and their number in  $[0, \pi/\sqrt{\beta}[$  is different from 2, then (4) has a  $T$ -periodic solution. In particular, this is true assuming that  $\Phi$  has a constant sign, which, in this context, corresponds to the so-called Landesman-Lazer condition. The choice of  $\nu = 3$  is crucial in [4], in order to exploit the isochronicity of the homogeneous equation.

The presence of the repulsive singularity has some analogies with the problem of a bouncing particle, which has been treated, e.g, in [3, 12, 13, 15]. For instance, concerning the simple model

$$\begin{cases} x'' + \beta x = e(t) \\ x(t_0) = 0 \end{cases} \Rightarrow x'(t_0^+) = -x'(t_0^-),$$

describing the behavior of a particle bouncing elastically against the barrier  $\{x = 0\}$ , with  $\beta > 0$  belonging to the set  $\Sigma$ , it was proved in [3] that, defining the function  $\Phi$  as above, the same type of result holds. In particular, as also shown in [15, Theorem 4.2], this is true if

$$\int_0^T e(t) |\sin(\sqrt{\beta}(t + \theta))| dt < 0, \quad \text{for every } \theta \in [0, T]. \quad (6)$$

The aim of this paper is to propose a kind of Landesman-Lazer condition in the case of a nonlinearity with a singularity like the one considered in [6], with the strict inequalities in (2) possibly replaced by equalities. We will succeed in replacing the upper inequality with an equality, by adding a suitable Landesman-Lazer type condition on that side. Moreover, condition (3) will be replaced by a one-sided control on the function  $g(t, x)$ . We point out that, under our assumptions, we are far from perturbing an isochronous oscillator like the one considered in [4].

Our situation seems to be more delicate than the one studied in [4], or, for the bouncing problem, in [3, 15]; for instance, dealing with equation (4) with  $\beta > 0$  belonging to  $\Sigma$ , and  $\nu \neq 3$ , it is not clear if a condition like (6) is sufficient for the existence of  $T$ -periodic solutions. In order to explain this, we appeal to the analogy between the perturbed isochronous Steen's equation and the bouncing problem. When looking for a priori estimates in the elastic bouncing problem, it is clear that the elasticity implies that all the bumps of the "limit function" must have the same height. Unfortunately, this behavior is not guaranteed for the limit function obtained for the singular periodic problem with lack of isochronicity (see Remark 2.5). Hence, since we are not able to predict that the height of each bump will be the same, we will need a Landesman-Lazer condition which refers to each bump, separately. Explicitly, defining

$$\psi(t) = \begin{cases} \sin(\sqrt{\beta}t) & \text{if } t \in \left[0, \frac{\pi}{\sqrt{\beta}}\right] \\ 0 & \text{if } t \in \left[\frac{\pi}{\sqrt{\beta}}, T\right], \end{cases}$$

instead of (6) we will ask a stronger condition, namely

$$\int_0^T e(t)\psi(t + \theta) dt < 0, \quad \text{for every } \theta \in [0, T]. \quad (7)$$

Under this assumption, we will prove that equation (4) has a  $T$ -periodic solution. However, the possibility of replacing condition (7) with (6) in our setting remains an open problem.

Notice that, following the arguments in [9], under the assumptions of Theorem 2.1 it is possible to prove that the radially symmetric system

$$\ddot{z} = -g(t, |z|) \frac{z}{|z|}, \quad z \in \mathbb{R}^M \setminus \{0\}$$

has infinitely many subharmonic solutions rotating around the origin. However, for brevity, we will not discuss this argument further on.

## 2 The main result

Let us consider the  $T$ -periodic problem

$$\begin{cases} x'' + g(t, x) = 0 \\ x(0) = x(T), x'(0) = x'(T), \end{cases} \quad (8)$$

where  $g : [0, T] \times ]0, +\infty[ \rightarrow \mathbb{R}$  is an  $L^\infty$ -Carathéodory function, namely

- $t \mapsto g(t, x)$  is measurable for every  $x > 0$ ;
- $x \mapsto g(t, x)$  is continuous, for almost every  $t \in [0, T]$ ;
- for every compact interval  $[a, b] \subset ]0, +\infty[$  there exists a constant  $C > 0$  such that

$$|g(t, x)| \leq C, \quad \text{for a.e. } t \in [0, T] \text{ and every } x \in [a, b].$$

Define, for a fixed nonnegative integer  $N$ ,

$$\mu_N = \left( \frac{N\pi}{T} \right)^2,$$

and set

$$\psi(t) = \begin{cases} \sin(\sqrt{\mu_{N+1}} t) & \text{if } t \in \left[0, \frac{T}{N+1}\right] \\ 0 & \text{if } t \in \left[\frac{T}{N+1}, T\right]. \end{cases}$$

Let us state our main result.

**Theorem 2.1.** *Assume the following hypotheses:*

(H1) *there exists a constant  $\hat{\mu}$  such that*

$$\liminf_{x \rightarrow +\infty} \frac{g(t, x)}{x} \geq \hat{\mu} > \mu_N,$$

*uniformly for almost every  $t \in [0, T]$ ;*

(H2) *there exists a constant  $\hat{\eta}$  such that*

$$g(t, x) \leq \mu_{N+1} x + \hat{\eta}, \quad \text{for a.e. } t \in [0, T] \text{ and every } x \geq 1.$$

*Moreover, for every  $\theta \in [0, T]$ ,*

$$\int_0^T \limsup_{x \rightarrow +\infty} (g(t, x) - \mu_{N+1} x) \psi(t + \theta) dt < 0; \quad (9)$$

(H3) there exist a constant  $\delta > 0$  and a continuous function  $f : ]0, \delta] \rightarrow \mathbb{R}$ , such that

$$g(t, x) \leq f(x) \quad \text{for a.e. } t \in [0, T] \text{ and every } x \in ]0, \delta],$$

with

$$\lim_{x \rightarrow 0^+} f(x) = -\infty, \quad \int_0^\delta f(x) dx = -\infty.$$

Then, problem (8) has a solution.

**Remark 2.2.** Theorem 2.1 clearly generalizes Theorem 1.1, since (2) implies (H1) and (H2), while (3) implies (H3), being in this case the left-hand side in (9) equal to  $-\infty$ .

Let us introduce, for  $\lambda \in [0, 1]$ , the family of functions

$$g_\lambda(t, x) = \lambda g(t, x) + (1 - \lambda) \left( -\frac{1}{x^3} + \frac{\mu_N + \mu_{N+1}}{2} x \right). \quad (10)$$

Moreover, for  $R > 1$ , let

$$\Lambda_R = \left\{ x \in C([0, T]) \mid \frac{1}{R} < x(t) < R \text{ for every } t \in [0, T] \right\}.$$

Following the arguments in [9, Section 2], it is sufficient to prove that there exists an  $R > 1$  such that, for any  $\lambda \in [0, 1]$ , every solution of the problem

$$\begin{cases} x'' + g_\lambda(t, x) = 0 \\ x(0) = x(T), x'(0) = x'(T) \end{cases} \quad (11)$$

belongs to  $\Lambda_R$ . We will prove this by means of the following two lemmas, the first one giving the estimate for the maxima and the second one focusing on minima.

Henceforth, when dealing with a solution  $x(t)$  of (11), in the Carathéodory sense, we will implicitly assume  $x(t) > 0$  for every  $t \in [0, T]$ . Moreover, we will assume that all the functions defined on  $[0, T]$  are extended by  $T$ -periodicity to the whole real line.

**Lemma 2.3.** *There exists a constant  $R > 1$  such that, for any  $\lambda \in [0, 1]$ , if  $x(t)$  is a solution of (11), then*

$$\max_{t \in [0, T]} x(t) < R.$$

*Proof.* By contradiction, assume that there exist  $(\lambda_n)_n \subset [0, 1]$  and a sequence  $(x_n)_n$  satisfying

$$\begin{cases} x_n'' + g_{\lambda_n}(t, x_n) = 0 \\ x_n(0) = x_n(T), x_n'(0) = x_n'(T), \end{cases} \quad (12)$$

with  $\|x_n\|_\infty \rightarrow +\infty$ . Up to a subsequence, we can assume  $\lambda_n \rightarrow \bar{\lambda} \in [0, 1]$ . We will prove the following claims.

**Claim 1** (estimate of minima). *There exists  $M > 1$  such that*

$$\min_{t \in [0, T]} x_n(t) \leq M,$$

for every  $n \in \mathbb{N}$ .

*Proof of Claim 1.* By contradiction, assume this is not the case, namely there exists a subsequence, still denoted by  $(x_n)_n$ , such that  $x_n(t) \rightarrow +\infty$  uniformly in  $t \in [0, T]$ . Integrating (12), we see that

$$\int_0^T g_{\lambda_n}(t, x_n(t)) dt = 0.$$

On the other hand, in view of (H1), there exists  $d > 0$  such that

$$g_{\lambda_n}(t, x) \geq \frac{\hat{\mu}}{2}x, \quad \text{for a.e. } t \in [0, T] \text{ and every } x \geq d.$$

Since, for  $n$  large,  $x_n(t) \geq d$  for every  $t \in [0, T]$ , this is clearly impossible.  $\square$

Claim 2. For  $n$  large,  $(x_n(t), x'_n(t))$  makes exactly  $N+1$  clockwise revolutions around the point  $(1, 0)$  when  $t$  varies from 0 to  $T$ .

*Proof of Claim 2.* We will apply some reasonings from [9], introducing, with the same notations, the function  $\mathcal{N} : ]0, +\infty[ \rightarrow \mathbb{R}$ , defined by

$$\mathcal{N}(u, v) = \left( \frac{1}{u^2} + u^2 + v^2 \right)^{\frac{1}{2}}.$$

Let  $\bar{\epsilon} > 0$  be such that

$$\bar{\epsilon} < \frac{T}{(N+1)(N+2)}. \quad (13)$$

It is possible to see (cf. [9, Lemma 2]) that there exist  $\zeta > 0$  and a sufficiently large  $R_1 > M$  with the following property:

if  $\tau_1 < \tau_2$  are such that  $(x_n(t), x'_n(t))$ , in the phase plane, makes exactly one rotation around the point  $(1, 0)$  when  $t$  varies from  $\tau_1$  to  $\tau_2$ , and  $\mathcal{N}(x_n(t), x'_n(t)) \geq R_1$  for every  $t \in [\tau_1, \tau_2]$ , then

$$\frac{T}{N+1} - \bar{\epsilon} \leq \tau_2 - \tau_1 \leq \zeta < \frac{T}{N}. \quad (14)$$

Choose now a positive integer  $\bar{n}$  such that

$$\bar{n} \left( \frac{T}{N+1} - \bar{\epsilon} \right) > T. \quad (15)$$

Using [9, Lemma 1], there exists  $R_2 > R_1$  such that, if, for some  $t_1 < t_2$ ,

$$\mathcal{N}(x_n(t_1), x'_n(t_1)) = R_1, \quad \mathcal{N}(x_n(t_2), x'_n(t_2)) = R_2,$$

and

$$R_1 < \mathcal{N}(x_n(t), x'_n(t)) < R_2 \quad \text{for every } t \in ]t_1, t_2[,$$

then  $(x_n(t), x'_n(t))$  makes, in the phase plane, at least  $\bar{n}$  clockwise turns around  $(1, 0)$  when  $t$  varies from  $t_1$  to  $t_2$ . Since  $\|x_n\|_\infty \rightarrow +\infty$ , for every  $n$  large we will have

$$\max\{\mathcal{N}(x_n(t), x'_n(t)) \mid t \in [0, T]\} > R_2.$$

Consequently,  $x_n(t)$  being  $T$ -periodic, for  $n$  large enough it will be  $\mathcal{N}(x_n(t), x'_n(t)) > R_1$  for every  $t \in [0, T]$ , otherwise  $(x_n(t), x'_n(t))$  would make at least  $\bar{n}$  turns around  $(1, 0)$

in the time  $T$ , which is impossible in view of (14) and (15).

Since  $\mathcal{N}(x_n(t), x'_n(t)) > R_1$  for every  $t \in [0, T]$ , enlarging  $R_1$ , if necessary, as shown in the proof of [9, Lemma 2],  $(x_n(t), x'_n(t))$  clockwise rotates at least once around  $(1, 0)$  in the time  $T$ . Taking into account (13) and (14), then, for  $n$  sufficiently large  $(x_n(t), x'_n(t))$  will make, in the phase plane, strictly more than  $N$  and strictly less than  $N + 2$  turns around  $(1, 0)$  in the time  $T$ . Since  $x_n(t)$  is  $T$ -periodic, this implies that  $(x_n(t), x'_n(t))$  will make exactly  $N + 1$  of such turns in the time  $T$ .  $\square$

Claim 3. *Setting*

$$v_n = \frac{x_n}{\|x_n\|_\infty},$$

*the sequence  $(v_n)_n$  converges uniformly, up to a subsequence, to a continuous function  $v$ . This function is not identically zero and it is twice continuously differentiable, except on a finite subset of  $[0, T]$ . Moreover,  $v$  satisfies  $v''(t) + \mu_{N+1}v(t) = 0$ , for almost every  $t \in [0, T]$ .*

*Proof of Claim 3.* For every  $n$ , the  $T$ -periodic function  $v_n$  solves

$$v_n'' + \frac{g_{\lambda_n}(t, x_n(t))}{\|x_n\|_\infty} v_n = 0. \quad (16)$$

Multiplying both sides of equation (16) by  $v_n$  and integrating between 0 and  $T$ , one has

$$\int_0^T v_n'(t)^2 dt = \int_0^T \frac{g_{\lambda_n}(t, x_n(t))}{\|x_n\|_\infty} v_n(t) dt.$$

By (H3), we can choose  $\bar{\delta} \in ]0, 1[$  such that, for every  $n$ ,

$$g_{\lambda_n}(t, x) \leq 0, \quad \text{for a.e. } t \in [0, T] \text{ and every } x \in ]0, \bar{\delta}].$$

For  $n$  large, it follows that

$$\begin{aligned} \int_0^T \frac{g_{\lambda_n}(t, x_n(t))}{\|x_n\|_\infty} v_n(t) dt &= \int_{\{x_n \leq \bar{\delta}\}} \cdots + \int_{\{\bar{\delta} < x_n < 1\}} \cdots + \int_{\{x_n \geq 1\}} \cdots \\ &\leq CT + \int_{\{x_n \geq 1\}} \frac{\mu_{N+1}x_n(t) + \hat{\eta}}{\|x_n\|_\infty} dt \leq (C + \mu_{N+1})T + 1, \end{aligned}$$

for a suitable constant  $C > 0$  coming from the Carathéodory condition. This immediately gives an estimate of the  $L^2$ -norm of  $v_n'$  which is independent of  $n$ . It follows that  $(v_n)_n$  is bounded in  $H^1(0, T)$ , so there exists  $v \in H^1(0, T)$  such that, up to a subsequence,  $v_n \rightharpoonup v$  (weakly) in  $H^1(0, T)$  and  $v_n \rightarrow v$  uniformly. Let us still denote such a subsequence by  $(v_n)_n$ . Since  $\|v_n\|_\infty = 1$  for every  $n$ , we have that  $\|v\|_\infty = 1$ . Hence,  $v$  is not identically zero; moreover, it is clear that, since  $v$  is the uniform limit of positive continuous  $T$ -periodic functions, it is a nonnegative continuous  $T$ -periodic function.

We now prove that  $v''(t) + \mu_{N+1}v(t) = 0$  almost everywhere in  $[0, T]$ . To this aim, define the open set

$$\Omega^+ = \{t \in \mathbb{R} \mid v(t) > 0\},$$

which is an at most countable union of open intervals, and consider a  $C^1$ -function  $\varphi$  with compact support  $K_\varphi$  contained in  $\Omega^+$ . Multiplying equation (16) by  $\varphi$  and integrating on  $K_\varphi$ , we obtain

$$\int_{K_\varphi} v_n'(t)\varphi'(t) dt = \int_{K_\varphi} \frac{g_{\lambda_n}(t, x_n(t))}{\|x_n\|_\infty} \varphi(t) dt = \int_{K_\varphi} \frac{g_{\lambda_n}(t, x_n(t))}{x_n(t)} v_n(t)\varphi(t) dt. \quad (17)$$

Observe now that, in view of the continuity of  $v$  (which is strictly greater than 0 on  $K_\varphi$ ) and recalling that  $K_\varphi$  is compact and strictly contained in  $\Omega^+$ , we have that  $x_n(t) \rightarrow +\infty$  uniformly for every  $t \in K_\varphi$ . By hypotheses (H1) and (H2), for every positive integer  $m$  there exists a sufficiently large  $n_m$  such that

$$\mu_N < \hat{\mu} - \frac{1}{m} \leq \frac{g_{\lambda_{n_m}}(t, x_{n_m}(t))}{x_{n_m}(t)} \leq \mu_{N+1} + \frac{1}{m}, \quad (18)$$

for almost every  $t \in K_\varphi$ . We can assume the sequence  $(n_m)_m$  to be strictly increasing. Considering the subsequence

$$\left( \frac{g_{\lambda_{n_m}}(t, x_{n_m}(t))}{x_{n_m}(t)} \right)_m,$$

it turns then out that it is bounded in  $L^2(K_\varphi)$ , so that (up to a further subsequence) it converges weakly in  $L^2(K_\varphi)$  to a function  $\gamma(t)$  which, in view of (18), satisfies

$$\mu_N < \hat{\mu} \leq \gamma(t) \leq \mu_{N+1}, \quad \text{for a.e. } t \in K_\varphi. \quad (19)$$

Since  $v_n \rightarrow v$  uniformly and  $v_n \rightharpoonup v$  in  $H^1(0, T)$ , passing to the limit in (17) yields then

$$\int_{K_\varphi} v'(t)\varphi'(t) dt = \int_{K_\varphi} \gamma(t)v(t)\varphi(t) dt.$$

By the uniqueness of the weak limit,  $\gamma(t)$  can be uniquely extended on the whole set  $\Omega^+$ , and hence we have

$$\int_{\Omega^+} v'(t)\varphi'(t) dt = \int_{\Omega^+} \gamma(t)v(t)\varphi(t) dt,$$

for every  $C^1$ -function  $\varphi$  with compact support in  $\Omega^+$ . This implies that  $v(t)$  is a *weak* solution of the equation

$$v'' + \gamma(t)v = 0 \quad (20)$$

in  $\Omega^+$ . Hence,  $v \in H_{\text{loc}}^2(\Omega^+)$ , and satisfies (20) for almost every  $t \in \Omega^+$ . Moreover,  $v(t)$  is continuously differentiable at any point  $t \in \Omega^+$ .

We now aim at proving that  $\gamma(t) = \mu_{N+1}$  for almost every  $t \in \Omega^+$ . Observe first that, writing the equation as a first order system, and taking into account that, for every  $n$ , the angular velocity of  $(v_n, v'_n)$  has to be nonpositive (see [9]), the set  $\{t \in [0, T] \mid x_n(t) = R_1\}$  (which is nonempty since  $R_1 > M$ ) is discrete. Moreover, coming back to Claim 2, we know that, in the phase plane,  $(x_n(t), x'_n(t))$  has to perform exactly  $N + 1$  turns around the point  $(1, 0)$  in the time  $T$ . Then, in the interval  $[0, 2T]$ , we can find

$$\alpha_1^n < \beta_1^n < \alpha_2^n < \beta_2^n < \dots < \alpha_{N+1}^n < \beta_{N+1}^n < \alpha_1^n + T$$

such that, setting  $\alpha_{N+2}^n = \alpha_1^n + T$ ,

$$x_n(t) > R_1 \quad \text{if } t \in ]\alpha_r^n, \beta_r^n[, r = 1, \dots, N + 1,$$

and

$$x_n(t) < R_1 \quad \text{if } t \in ]\beta_r^n, \alpha_{r+1}^n[, r = 1, \dots, N + 1.$$

Since both the sequences  $(\beta_r^n)_n$  and  $(\alpha_r^n)_n$  are bounded, there exist  $\xi_r^-$  and  $\xi_r^+$ , with  $r = 1, \dots, N + 1$ , such that, up to subsequences,

$$\alpha_r^n \rightarrow \xi_r^-, \quad \beta_r^n \rightarrow \xi_r^+, \quad r = 1, \dots, N + 1,$$

with

$$\xi_1^- \leq \xi_1^+ \leq \xi_2^- \leq \xi_2^+ \leq \dots \leq \xi_{N+1}^- \leq \xi_{N+1}^+ \leq \xi_1^- + T. \quad (21)$$

By the estimates in [9, Lemma 2], given any  $\epsilon > 0$ , for  $n$  sufficiently large we have

$$\beta_r^n - \alpha_r^n \geq \frac{T}{N + 1} - \epsilon, \quad r = 1, \dots, N + 1.$$

Passing to the limit, being  $\epsilon$  arbitrary, we deduce

$$\xi_r^+ - \xi_r^- \geq \frac{T}{N + 1}, \quad r = 1, \dots, N + 1.$$

Since, by (21),  $\xi_{N+1}^+ - \xi_1^- \leq T$ , this implies

$$\xi_r^+ = \xi_r^- + \frac{T}{N + 1}, \quad \text{and} \quad \xi_{r+1}^- = \xi_r^+, \quad r = 1, \dots, N + 1.$$

For simplicity of notations, we set  $\xi_r = \xi_r^-$ , for  $r = 1, \dots, N + 1$ . Recalling that

$$v_n(\alpha_r^n) = v_n(\beta_r^n) = \frac{R_1}{\|x_n\|_\infty},$$

we deduce that  $v(\xi_r) = 0$  for every  $r = 1, \dots, N + 1$ , since  $v$  is continuous and  $v_n \rightarrow v$  uniformly.

Let us now focus on an interval  $[\tilde{\alpha}, \tilde{\beta}]$  such that  $v > 0$  on  $] \tilde{\alpha}, \tilde{\beta} [$ , with  $v(\tilde{\alpha}) = v(\tilde{\beta}) = 0$ . Since, on  $] \tilde{\alpha}, \tilde{\beta} [$ ,  $v$  is continuously differentiable and satisfies (20) almost everywhere, writing

$$v(t) = \rho(t) \cos(\theta(t)), \quad v'(t) = \rho(t) \sin(\theta(t)),$$

we have

$$-\theta'(t) = \frac{\gamma(t)v(t)^2 + v'(t)^2}{v(t)^2 + v'(t)^2},$$

which yields, in view of (19),

$$\frac{-\theta'(t)}{\mu_{N+1} \cos^2 \theta(t) + \sin^2 \theta(t)} \leq 1 < \frac{-\theta'(t)}{\hat{\mu} \cos^2 \theta(t) + \sin^2 \theta(t)}.$$

Consequently, integrating between  $\tilde{\alpha}$  and  $\tilde{\beta}$ , we infer, taking into account that  $\hat{\mu} > \mu_N$ ,

$$\frac{T}{N + 1} \leq \tilde{\beta} - \tilde{\alpha} < \frac{T}{N}.$$

Since, however,  $v(\xi_r) = 0$  for every  $r = 1, \dots, N + 1$ , and the points  $\xi_r$  are equally distributed at a distance  $\frac{T}{N+1}$  one from the other, necessarily  $\tilde{\beta} - \tilde{\alpha} = \frac{T}{N+1}$ . This means that, whenever  $v$  becomes positive, it is forced to remain positive for a time exactly equal to  $\frac{T}{N+1}$ . Recalling that  $v$  solves  $v'' + \gamma(t)v = 0$  on  $] \tilde{\alpha}, \tilde{\beta} [$ , we now pass to generalized polar coordinates by writing

$$v(t) = \frac{1}{\sqrt{\mu_{N+1}}} \hat{\rho}(t) \cos(\hat{\theta}(t)), \quad v'(t) = \hat{\rho}(t) \sin(\hat{\theta}(t)).$$

We have, integrating  $\hat{\theta}'(t)$  on  $[\tilde{\alpha}, \tilde{\beta}]$ ,

$$\pi = \sqrt{\mu_{N+1}} \int_{\tilde{\alpha}}^{\tilde{\beta}} \frac{\gamma(t)v(t)^2 + v'(t)^2}{\mu_{N+1}v(t)^2 + v'(t)^2} dt \leq \sqrt{\mu_{N+1}} \frac{T}{N+1} = \pi,$$

from which we deduce that

$$\gamma(t) = \mu_{N+1}, \quad \text{for a.e. } t \in [\tilde{\alpha}, \tilde{\beta}].$$

This means, in particular, that  $\Omega^+$  is the union of some intervals of the type  $]\xi_r, \xi_{r+1}[$ , on which we have

$$v(t) = c_r \sin(\sqrt{\mu_{N+1}}(t - \xi_r)),$$

where, for every  $r$ , the constants  $c_r$  are in  $[0, 1]$ , and at least one of them is equal to 1. The claim is thus proved.  $\square$

### Conclusion of the proof of Lemma 2.3.

Let us focus again on an interval  $[\tilde{\alpha}, \tilde{\beta}]$  such that  $v(\tilde{\alpha}) = v(\tilde{\beta}) = 0$ , and  $v > 0$  on  $]\tilde{\alpha}, \tilde{\beta}[$ . Notice first that, since  $\gamma(t) = \mu_{N+1}$  almost everywhere in  $\Omega^+$ , necessarily it will be  $\lambda_n \rightarrow 1$ . Indeed, if it were  $\lambda_n \rightarrow \bar{\lambda} < 1$ , taking any interval  $I \subset ]\tilde{\alpha}, \tilde{\beta}[$ , for every  $t \in I$  it would be, for  $n$  large,

$$\frac{g_{\lambda_n}(t, x_n(t))}{x_n(t)} \leq \tilde{\zeta} < \mu_{N+1},$$

for a suitable constant  $\tilde{\zeta}$ , and, taking the weak limit, we would reach the contradiction  $\gamma(t) < \mu_{N+1}$ , for almost every  $t \in I$ . By the previous discussion, we know that  $\tilde{\beta} - \tilde{\alpha} = \frac{T}{N+1}$  and that, on  $]\tilde{\alpha}, \tilde{\beta}[$ ,  $v$  solves the Dirichlet problem

$$\begin{cases} v'' + \mu_{N+1}v = 0 \\ v(\tilde{\alpha}) = 0 = v(\tilde{\beta}). \end{cases}$$

Explicitly, for some  $C > 0$ ,

$$v(t) = C \sin\left(\frac{(N+1)\pi}{T}(t - \tilde{\alpha})\right). \quad (22)$$

Let  $(\phi_k)_{k \geq 1}$  be the orthonormal  $L^2(\tilde{\alpha}, \tilde{\beta})$ -basis made of the solutions of the Dirichlet problem

$$\begin{cases} \phi_k'' + \mu_k \phi_k = 0 \\ \phi_k(\tilde{\alpha}) = 0 = \phi_k(\tilde{\beta}), \end{cases}$$

where  $\mu_k$  is the  $k$ -th eigenvalue. Let us denote by  $\langle \cdot, \cdot \rangle$  the scalar product in  $L^2(\tilde{\alpha}, \tilde{\beta})$  and by  $\|\cdot\|_2$  the  $L^2(\tilde{\alpha}, \tilde{\beta})$ -norm. We write, for every  $n$ , the Fourier series of  $x_n$ ,

$$x_n = \sum_{k=1}^{+\infty} \langle x_n, \phi_k \rangle \phi_k.$$

It is useful to split such an expression as follows:

$$x_n = x_n^0 + x_n^\perp,$$

where

$$x_n^0 = \langle x_n, \phi_{N+1} \rangle \phi_{N+1},$$

while

$$x_n^\perp = \sum_{k \neq N+1} \langle x_n, \phi_k \rangle \phi_k.$$

It is known that  $x_n^\perp$  is  $L^2$ -orthogonal to  $x_n^0$  in view of the properties of the eigenfunctions  $\phi_k$ . Dividing  $x_n$  by  $\|x_n\|_\infty$ , we have  $v_n = v_n^0 + v_n^\perp$ , with

$$v_n^0 = \frac{x_n^0}{\|x_n\|_\infty}, \quad v_n^\perp = \frac{x_n^\perp}{\|x_n\|_\infty}.$$

Since  $v = v^0$ , we have

$$v_n^0 = \frac{\langle v_n, v \rangle}{\|v\|_2} \rightarrow v, \quad \text{uniformly in } t \in [\tilde{\alpha}, \tilde{\beta}].$$

As

$$x_n'' = \sum_{k=1}^{+\infty} \langle x_n, \phi_k \rangle \phi_k'' = - \sum_{k=1}^{+\infty} \langle x_n, \phi_k \rangle \mu_k \phi_k,$$

we have that

$$(x_n'')^0 = (x_n^0)'', \quad (x_n'')^\perp = (x_n^\perp)''.$$

Multiplying equation (12) by  $v_n^0$  and integrating between  $\tilde{\alpha}$  and  $\tilde{\beta}$  we then have

$$- \int_{\tilde{\alpha}}^{\tilde{\beta}} (x_n^0)''(t) v_n^0(t) dt = \int_{\tilde{\alpha}}^{\tilde{\beta}} g_{\lambda_n}(t, x_n(t)) v_n^0(t) dt. \quad (23)$$

Integrating twice by parts, we obtain

$$\int_{\tilde{\alpha}}^{\tilde{\beta}} (x_n^0)''(t) v_n^0(t) dt = \int_{\tilde{\alpha}}^{\tilde{\beta}} x_n^0(t) (v_n^0)''(t) dt = - \int_{\tilde{\alpha}}^{\tilde{\beta}} \mu_{N+1} x_n^0(t) v_n^0(t) dt. \quad (24)$$

Notice that, by (H3), there exists  $\omega \in ]0, \min\{\delta, 1\}[$  such that

$$g(t, x) \leq f(x) \leq 0, \quad \text{for every } x \in ]0, \omega[.$$

Define the two  $L^\infty$ -Carathéodory functions

$$\sigma(t, x) = \begin{cases} g(t, x) & \text{if } x \in ]0, \frac{\omega}{2}[ \\ \frac{2}{\omega} g(t, x)(\omega - x) & \text{if } x \in [\frac{\omega}{2}, \omega] \\ 0 & \text{if } x \in ]\omega, +\infty[, \end{cases}$$

and

$$r(t, x) = g(t, x) - \mu_{N+1} x - \sigma(t, x).$$

Using (H2), together with the fact that  $\sigma(t, x) \leq 0$ , there exists a positive constant  $\tilde{\eta}$  such that

$$r(t, x) \leq \tilde{\eta}, \quad \text{for a.e. } t \in [0, T], \text{ and every } x > 0. \quad (25)$$

As a consequence,

$$\begin{aligned}
\int_{\tilde{\alpha}}^{\tilde{\beta}} g_{\lambda_n}(t, x_n(t))v_n^0(t) dt &= \lambda_n \int_{\tilde{\alpha}}^{\tilde{\beta}} \sigma(t, x_n(t))v_n^0(t) dt + \lambda_n \int_{\tilde{\alpha}}^{\tilde{\beta}} \mu_{N+1}x_n(t)v_n^0(t) dt + \\
&+ \lambda_n \int_{\tilde{\alpha}}^{\tilde{\beta}} r(t, x_n(t))v_n^0(t) dt + (1 - \lambda_n) \int_{\tilde{\alpha}}^{\tilde{\beta}} \left( -\frac{1}{x_n(t)^3} + \frac{\mu_N + \mu_{N+1}}{2}x_n(t) \right) v_n^0(t) dt \\
&\leq \int_{\tilde{\alpha}}^{\tilde{\beta}} \mu_{N+1}x_n^0(t)v_n^0(t) dt + \lambda_n \int_{\tilde{\alpha}}^{\tilde{\beta}} r(t, x_n(t))v_n^0(t) dt,
\end{aligned}$$

where here we used the fact that  $x_n$  and  $v_n^0$  are positive. Using (23), (24) and the fact that  $\lambda_n \rightarrow 1$ , we deduce that, for  $n$  sufficiently large,

$$\int_{\tilde{\alpha}}^{\tilde{\beta}} r(t, x_n(t))v_n^0(t) dt \geq 0.$$

As  $v_n^0 \rightarrow v$  uniformly, by (25) the integrand is bounded from above by a positive constant. It is then possible to apply Fatou's lemma, so that

$$\int_{\tilde{\alpha}}^{\tilde{\beta}} \limsup_{n \rightarrow +\infty} r(t, x_n(t))v_n^0(t) dt \geq 0,$$

whence, since  $\lim_{n \rightarrow +\infty} v_n^0(t) = v(t)$  for every  $t \in [\tilde{\alpha}, \tilde{\beta}]$ ,

$$\int_{\tilde{\alpha}}^{\tilde{\beta}} \limsup_{n \rightarrow +\infty} r(t, x_n(t))v(t) dt \geq 0.$$

As

$$r(t, x) = g(t, x) - \mu_{N+1}x, \quad \text{for a.e. } t \in [0, T] \text{ and every } x \geq 1,$$

and  $x_n(t) \rightarrow +\infty$  for every  $t \in ]\tilde{\alpha}, \tilde{\beta}[$ , we deduce that

$$\int_{\tilde{\alpha}}^{\tilde{\beta}} \limsup_{x \rightarrow +\infty} (g(t, x) - \mu_{N+1}x)v(t) dt \geq 0.$$

Since, in view of (22),  $v(t) = C\psi(t - \tilde{\alpha})$  for every  $t \in [\tilde{\alpha}, \tilde{\beta}]$ , this contradicts (9).  $\square$

We now prove the counterpart of Lemma 2.3 for minima.

**Lemma 2.4.** *There exists a constant  $R' \in ]0, 1[$  such that, for any  $\lambda \in [0, 1]$ , if  $x(t)$  is a solution of (11), then*

$$\min_{t \in [0, T]} x(t) > R'.$$

*Proof.* By contradiction, assume that there exist two sequences  $(\lambda_n)_n \subset [0, 1]$  and  $(x_n)_n$ , with  $x_n(t)$  solving (11) for  $\lambda = \lambda_n$ , such that  $\min_{t \in [0, T]} x_n(t) \rightarrow 0$ . Let  $R > 1$  be as in Lemma 2.3 and fix  $\epsilon > 0$  sufficiently small. As we have already recalled, by [9, Lemma 2] there exists a sufficiently large  $R_1 > \mathcal{N}(R, 0)$  such that, if  $(x_n(t), x'_n(t))$  makes exactly one rotation around the point  $(1, 0)$  when  $t$  varies from  $\tau_1$  to  $\tau_2$ , and  $\mathcal{N}(x_n(t), x'_n(t)) \geq R_1$  for every  $t \in [\tau_1, \tau_2]$ , then

$$\tau_2 - \tau_1 \geq \frac{T}{N+1} - \epsilon.$$

Choose now a positive integer  $\bar{n}$  such that

$$\bar{n} \left( \frac{T}{N+1} - \epsilon \right) > T.$$

Recalling again [9, Lemma 1], as used in the proof of Lemma 2.3, there exists  $R_2 > R_1$  such that if

$$\mathcal{N}(x_n(t_1), x'_n(t_1)) = R_1, \quad \mathcal{N}(x_n(t_2), x'_n(t_2)) = R_2,$$

for some  $t_1 < t_2$ , and

$$R_1 < \mathcal{N}(x_n(t), x'_n(t)) < R_2 \quad \text{for every } t \in ]t_1, t_2[,$$

then  $(x_n(t), x'_n(t))$  turns at least  $\bar{n}$  times around  $(1, 0)$  in the phase plane when  $t$  varies from  $t_1$  to  $t_2$ . As  $\min_{t \in [0, T]} x_n(t) \rightarrow 0$ , for every  $n$  large we will have

$$\max\{\mathcal{N}(x_n(t), x'_n(t)) \mid t \in [0, T]\} > R_2.$$

Since  $x_n(t)$  is  $T$ -periodic, and in view of the continuity of  $\mathcal{N}(x_n(t), x'_n(t))$ , for  $n$  large enough it follows that  $\mathcal{N}(x_n(t), x'_n(t)) > R_1$  for every  $t \in [0, T]$ , otherwise  $(x_n(t), x'_n(t))$  would turn at least  $\bar{n}$  times around  $(1, 0)$  in the time  $T$ , which is impossible. Choosing  $\bar{t}_n$  such that  $x_n(\bar{t}_n) = \max_{t \in [0, T]} x_n(t)$ , being  $x'_n(\bar{t}_n) = 0$  it would then be, for  $n$  large,

$$\mathcal{N}(x_n(\bar{t}_n), 0) > R_1,$$

which, since  $x_n(\bar{t}_n) < R$ , contradicts the inequality  $\mathcal{N}(R, 0) < R_1$ .  $\square$

**Remark 2.5.** As we mentioned in the Introduction, in the special case of equation (4), condition (9) reads as (7). Let us try to explain why we are not able to replace (7) with the more general assumption (6). We recall that, in [9], a curve  $\Gamma : [0, +\infty[ \rightarrow \mathbb{R}^2$  was constructed in order to control the solutions in the phase plane. One could try to estimate the height of the bumps of  $v(t)$  by using such a curve for the solutions  $x_n(t)$ . However, consider for example equation (5) and define  $g_\lambda(t, x)$  as in (10). We would then have

$$g_\lambda(t, x) = -\frac{1}{x^3} + \left( \frac{\mu_N + \mu_{N+1}}{2} + \lambda \frac{\mu_{N+1} - \mu_N}{2} \right) x.$$

Hence, the curve  $\Gamma(t)$ , as it is constructed in [9], should use the level lines of the functions

$$V_1(u, v) = \frac{1}{2} \left( \frac{1}{u^2} + \mu_{N+1} u^2 + v^2 \right), \quad V_2(u, v) = \frac{1}{2} \left( \frac{1}{u^2} + \frac{\mu_N + \mu_{N+1}}{2} u^2 + v^2 \right).$$

A straightforward computation shows, however, that, denoting by  $P_m = (u_m, 0)$  and  $P_{m+1} = (u_{m+1}, 0)$  two consecutive intersections of  $\Gamma(t)$  with the half-line  $\{(u, v) \in \mathbb{R}^2 \mid u \geq 1, v = 0\}$ , it is

$$\liminf_{m \rightarrow +\infty} \frac{u_{m+1}}{u_m} > 1.$$

This fact makes us unable to prove that the height of the bumps of  $v(t)$ , which is obtained as the limit of the normalized sequence  $(v_n)_n$ , is always the same. This is why we did not manage, in our setting, to assume a Landesman-Lazer condition like (6).

## References

- [1] S. Bolotin and R.S. Mackay, *Isochronous potentials*, in: Localization and Energy Transfer in Nonlinear Systems, World Sci, 2003, 217–224.
- [2] D. Bonheure and C. De Coster, *Forced singular oscillators and the method of lower and upper solutions*, Topol. Meth. Nonlin. Anal. **22** (2003), 297–317.
- [3] D. Bonheure and C. Fabry, *Periodic motions in impact oscillators with perfectly elastic bounces*, Nonlinearity **15** (2002), 1281–1297.
- [4] D. Bonheure, C. Fabry and D. Smets, *Periodic solutions of forced isochronous oscillators at resonance*, Discrete Contin. Dyn. Syst. **8** (2002), 907–930.
- [5] M. del Pino and R. Manásevich, *Infinitely many  $T$ -periodic solutions for a problem arising in nonlinear elasticity*, J. Differential Equations **103** (1993), 260–277.
- [6] M. del Pino, R. Manásevich and A. Montero,  *$T$ -periodic solutions for some second order differential equations with singularities*, Proc. Royal Soc. Edinburgh **120A** (1992), 231–243.
- [7] A. Fonda, *Periodic solutions of scalar second order differential equations with a singularity*, Académie Royale de Belgique, Brussels, 1993.
- [8] A. Fonda, R. Manásevich, and F. Zanolin, *Subharmonic solutions for some second-order differential equations with singularities*, SIAM J. Math. Anal. **24** (1993), 1294–1311.
- [9] A. Fonda and R. Toader, *Radially symmetric systems with a singularity and asymptotically linear growth*, Nonlinear Anal., to appear.
- [10] S. Fučík, Solvability of Nonlinear Equations and Boundary Value Problems, Reidel, Boston, 1980.
- [11] P. Habets and L. Sanchez, *Periodic solutions of some Liénard equations with singularities*, Proc. Amer. Math. Soc. **109** (1990), 1035–1044.
- [12] M.-Y. Jiang, *Periodic solutions of second order differential equations with an obstacle*, Nonlinearity **19** (2006), 1165–1183.
- [13] A. C. Lazer and P. J. McKenna, *Periodic bouncing for a forced linear spring with obstacle*, Differential Integral Equations **5** (1992), 165–172.
- [14] A. C. Lazer and S. Solimini, *On periodic solutions of nonlinear differential equations with singularities*, Proc. Amer. Math. Soc. **99** (1987), 109–114.
- [15] R. Ortega, *Dynamics of a forced oscillator having an obstacle*, in: Variational and Topological Methods in the Study of Nonlinear Phenomena (Pisa, 2000), 75–87, Progr. Nonlinear Differential Equations Appl., 49, Birkhäuser, Boston, 2002.
- [16] R. Ortega, *Periodic perturbations of an isochronous center*, Qual. Theory Dyn. Syst. **3** (2002), 83–91.

- [17] I. Rachůnková, S. Staněk and M. Tvrdý, *Solvability of Nonlinear Singular Problems for Ordinary Differential Equations*, Hindawi, New York, 2008.
- [18] R. Redheffer, *Steen's equation and its generalizations*, *Aequationes Math.* **58** (1999), 60–72.
- [19] Z. Wang, *Periodic solutions of the second-order differential equations with singularity*, *Nonlinear Anal.* **58** (2004), 319–331.
- [20] J. Xia and Z. Wang, *Existence and multiplicity of periodic solutions for the Duffing equation with singularity*, *Proc. Royal Soc. Edinburgh* **137A** (2007), 1–21.
- [21] J. Xia, Z. Wang and D. Zheng, *Periodic solutions of Duffing equations with semi-quadratic potential and singularity*, *J. Math. Anal. Appl.* **321** (2006), 273–285.
- [22] M. Zhang, *A relationship between the periodic and the Dirichlet BVPs of singular differential equations*, *Proc. Royal Soc. Edinburgh* **128A** (1998), 1099–1114.
- [23] M. Zhang, *Periodic solutions of equations of Emarkov-Pinney type*, *Adv. Nonlinear Stud.* **6** (2006), 57–67.

Authors' address:

Alessandro Fonda  
 Dipartimento di Matematica e Informatica  
 Università di Trieste  
 P.le Europa 1  
 I-34127 Trieste  
 Italy  
 e-mail: a.fonda@units.it

Maurizio Garrione  
 SISSA - International School for Advanced Studies  
 Via Bonomea 265  
 I-34136 Trieste  
 Italy  
 e-mail: garrione@sissa.it

Mathematics Subject Classification: 34C25; 34B16.

Keywords: periodic solutions; repulsive singularity; resonance; Landesman-Lazer condition.